

NET WEIGHT 50 LBS  
U L T I M A T E S H O W  
G O A T P E L L E T

MEDICATED

For the prevention of coccidiosis in young goats caused by *Eimeria cristenseni* and *Eimeria ninakohylyakimovae*. Feed at least 28 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard.

ACTIVE DRUG INGREDIENT

Decoquinate.....13.6g/ton

.....GUARANTEED ANALYSIS.....

CRUDE PROTEIN.....NOT LESS THAN.....17.38%  
CRUDE FAT.....NOT LESS THAN..... 3.20%  
CRUDE FIBER.....NOT MORE THAN.....17.20%  
ASH.....NOT MORE THAN..... 8.90%

.....

LIST OF INGREDIENTS

Sun-Cured Alfalfa Meal, Ground Corn, Ground Oats, Soybean Meal, Soyhull Pellets (7.5%), Dried Distillers Grains w/Solubles, Cane Molasses, Fish Meal, Ammonium Chloride, Canola Oil, Calcium Carbonate, Salt (Sodium Chloride), Dried *Saccharomyces cerevisiae* (Diamond V XP Yeast), Sodium Sesquicarbonate, Mono Dicalcium Phosphate, Vitamin A Acetate, Vitamin D3 Supplement, Vitamin E, Manganous Oxide, Zinc Sulfate, Ferrous Sulfate, Magnesium Sulfate, Potassium Sulfate, Ethylenediamine Dihydriodide, Cobalt Sulfate, Sodium Selenite.

WARNING: Do not feed to goats producing milk for food.